

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 207/08, 211/10, A61K 31/40, 31/445	A1	(11) International Publication Number: WO 00/27811 (43) International Publication Date: 18 May 2000 (18.05.00)
(21) International Application Number: PCT/US99/26798 (22) International Filing Date: 9 November 1999 (09.11.99) (30) Priority Data: 60/108,060 12 November 1998 (12.11.98) US (71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US). (72) Inventors: DUBOWCHIK, Gene, M.; 65 Spring Street, Middlefield, CT 06455 (US). DITTA, Jonathan, L.; 6111 Town Place, Middletown, CT 06457 (US). PROVENCAL, David, P.; 131 Trolley Crossing Lane, Middletown, CT 06457 (US). DENHART, Derek, J.; 188 N. Whittlesey Avenue, Wallingford, CT 06492 (US). (74) Agent: DUBOFF, Samuel, J.; Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: NEUROTROPHIC DIAMIDE AND CARBAMATE AGENTS		
(57) Abstract The present invention relates to the design, synthesis, and the peptidyl-prolyl isomerase (PPIase or rotamase) inhibitory activity of novel pyrrolidinemethyl diamide and carbamate compounds that are neurotrophic agents (i.e. compounds capable of stimulating growth or proliferation of nervous tissue) and that bind to immunophilins such as FKBP12 and inhibit their rotamase activity. This invention also relates to pharmaceutical compositions comprising these compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NEUROTROPHIC DIAMIDE AND CARBAMATE AGENTS**CROSS-REFERENCE TO RELATED APPLICATIONS**

- 5 This application claims the priority of U.S. Provisional Application Serial No. 60/108,060 filed November 12, 1998.

BACKGROUND OF THE INVENTION

- 10 Immunophilins are cytosolic proteins that possess peptidyl-prolyl-*cis-trans* isomerase (PPIase or rotamase) activity. This family of proteins behave as chaperone molecules causing *cis-trans* isomerization of peptide-prolyl bonds that could be a rate limiting step in the correct folding of certain proteins. They are also involved in many cellular signal transduction pathways as partners in multiprotein complexes for which
- 15 binding in the rotamase active site, but not rotamase activity *per se*, appears to be important (Rühlmann, et al., *Immunobiol.*, 198, pp. 192-206 (1998)). Immunosuppressive drugs such as FK506, rapamycin and cyclosporin A bind to specific groups of immunophilins. FK506 and rapamycin bind to the so-called FK506-binding proteins (FKBPs), whereas the cyclophilins binds to cyclosporin A. It
- 20 has been shown that binding to the 12kD immunophilin FKBP12 is necessary for FK506 to elicit its immunosuppressive activity. Subsequently, it was also found that FK506 has two binding domains: one that binds to FKBP12 and the other (the effector domain) for the complex of FK506 and FKBP12 that binds to the serine/threonine phosphatase, calcineurin. This complexation inhibits calcineurin and prevents the
- 25 proliferation of T-lymphocytes (i.e. immunosuppression). Rapamycin has an effector domain of a different structure, and its complex with FKBP12 binds to a different target protein that, also results in immunosuppression. For a review, see S.L. Schreiber, et al., *Tetrahedron*, 48, pp. 2545-2558 (1992). Some of the other proteins

with which FKBP12 is known to interact include the TGF β receptor I (Wang, et al.,
Science, 265, pp. 674-676 (1994)), the IP₃ receptor and the ryanodine receptor
(Cameron, et al., *J. Biol. Chem.*, 272, pp. 27582-27588 (1997)). In the case of the
TGF β system, it has been suggested that FKBP12 binding inhibits unregulated
5 signalling with consequences for differentiation, apoptosis and proliferation (Wang, et
al., *Cell*, 86, pp. 435-444 (1996)).

While FK506 exhibits immunosuppressive effects, analogs lacking the
calcineurin binding effector domain are devoid of immunosuppressive activity. Many
10 small molecules that contain the essential elements of the FKBP12 binding domain of
FK506 but lack the calcineurin binding domain were found to retain high affinity
binding to FKBP12, and behave as rotamase inhibitors (D.S. Yamshita, et al., *Bioorg.
Med. Chem. Lett.*, 4, pp. 325-328 (1994); D.M. Armistead, et al., *Acta Cryst. D*, 51,
pp. 522-528 (1995)).

15

FK506 has been shown to possess neurotrophic properties *in vitro* and *in vivo*
(W.E. Lyons, et al., *Proc. Natl. Acad. Sci USA*, 91, pp. 3191-3195 (1994); B.G.
Gold, et al., *J. Neurosci.*, 15, pp. 7509-7516 (1995)). However, its
immunosuppressive properties as well as other serious side effects are drawbacks to its
20 use as a neuroregenerative agent. Recently, *in vitro* studies in PC12 cells, SY5Y cells,
and chick sensory dorsal root ganglion explant cultures have shown that small
molecule, nonimmunosuppressive FKBP12 rotamase inhibitors also promote neurite
outgrowth, and a number of these compounds have shown utility in reversal of CNS
lesioning and nerve crush in animal models (G.S. Hamilton, et al., *Curr. Pharm.*
25 *Design*, 3, pp. 405-428 (1997); B.G. Gold, et al., *Exp. Neurol.*, 147, pp. 269-278
(1997)). Thus, while the calcineurin binding domain of FK506 is necessary for
immunosuppressive activity, it is not required for neurotrophic activity.

A 10-50 fold elevated expression of immunophilins in the central nervous system in comparison with the immune system is well documented (S.H. Snyder, et al., *Nature Med.*, 1, pp. 32-37 (1995)). Recently, augmented expression of FKBP12 m-RNA following facial nerve crush and sciatic nerve lesions was established in facial and lumbar motor neurons. The observed augmentation paralleled the enhanced expression of growth associated protein GAP43 mRNA (B.G. Gold, et al., *Neurosci. Lett.*, 241, pp. 25-28 (1998)). These observations make FKBP12 an attractive target for developing nonimmunosuppressive rotamase inhibitors which promote neurite outgrowth. Such compounds are potential therapeutics to reverse neuronal damage caused by neurodegenerative disease or physical trauma.

There have been disclosures of related compounds for overcoming multidrug resistance (MDR) or as immunosuppressants such as:

WO 94/07858 published 4/14/94
WO 92/19593 published 11/12/92
U.S. Patent 5,622,970 granted 4/22/97
U.S. Patent 5,330,993 granted 7/19/94
U.S. Patent 5,192,773 granted 3/9/93
U.S. Patent 5,516,797 granted 5/14/96
WO 92/21313 published 12/10/92
European Application 564924 published 10/13/93
European Application 405994 published 1/2/91

Other prior art disclosing related compounds having neurotrophic activity are:

WO 96/40140 published 12/19/96
WO 96/40633 published 12/19/96

- WO 97/16190 published 5/9/97
WO 96/41609 published 12/27/96
U.S. Patent 5,696,135 granted 12/9/97
WO 97/36869 published 10/9/97
5 U.S. Patent 5,721,256 granted 2/24/98
U.S. Patent 5,654,332 granted 8/5/97
WO 98/13343 published 4/2/98
WO 98/13355 published 4/2/98
WO 98/20891 published 5/22/98
10 WO 98/20892 published 5/22/98
WO 98/20893 published 5/22/98
WO 98/29116 published 7/9/98
WO 98/29117 published 7/9/98
U.S. Patent 5,780,484 granted 7/14/98
15 U.S. Patent 5,786,378 granted 7/28/98
U.S. Patent 5,795,908 granted 8/18/98
U.S. Patent 5,798,355 granted 8/25/98
U.S. Patent 5,801,187 granted 9/1/98
U.S. Patent 5,801,197 granted 9/1/98

20

Since there are relatively few FKBP12-binding compounds that are known to stimulate neurite growth, there remains a great need for additional neurotrophic, FKBP12-binding compounds.

25

SUMMARY OF THE INVENTION


Surprisingly, applicant has solved the aforementioned problem. The present invention relates to novel diamide and carbamate compounds and pharmaceutical compositions thereof that possess neurotrophic properties.

DETAILED DESCRIPTION OF THE INVENTION

According to one embodiment, the present invention provides:

10

A compound with affinity for an FK506 binding protein having the formula (I):



15 and pharmaceutically acceptable salts thereof, wherein:

Z is O, NH, N(C₁-C₃)-alkyl or CH₂;

X is O or F₂;

20

B and C are independently hydrogen, (C₁-C₄)-alkyl, or benzyl;

J is hydrogen, (C₁-C₄)-alkyl or benzyl;

K is (C₁-C₄)-straight or branched alkyl, benzyl or cyclohexylmethyl, or wherein J and K may be taken together to form a 4-7 membered heterocyclic ring which may contain a heteroatom selected from the group consisting of O, S, SO or SO₂, N, and NR, wherein R is selected from the group consisting of hydrogen, (C₁-C₄)-straight or
5 branched alkyl, (C₂-C₄)-straight or branched alkenyl or alkynyl, and (C₁-C₄)-bridging alkyl wherein a bridge is formed between the nitrogen and a carbon atom of said heteroatom-containing chain to form a ring, and wherein said ring is optionally fused to an Ar group;

10 wherein the stereochemistry at carbon position 1 is R or S;

D is (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl, (C₅-C₇)-cycloalkyl or (C₅-C₇)-cycloalkenyl substituted with (C₁-C₄)-straight or branched alkyl or (C₂-C₄)-straight or branched alkenyl, O-(C₁-C₄)-straight or branched alkyl, O-(C₂-C₄)-straight or branched alkenyl, 2-indolyl, 3-indolyl,
15 [(C₁-C₄)-alkyl or (C₂-C₄)-alkenyl]-Ar or Ar;

Ar is a carbocyclic aromatic group selected from the group consisting of phenyl, 1-naphthyl, 2-naphthyl, indenyl, azulenyl, fluorenyl, and anthracenyl; or a heterocyclic aromatic group selected from the group consisting of 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl,
20 pyrazolyl, isoxazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,3-triazolyl, 1,3,4-thiadiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-triazinyl, 1,3,5-trithianyl, indoliziny, indolyl, isoindolyl, 3H-indolyl, indolinyl, benzo[b]furanyl, benzo[b]thiophenyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, 4H-quinoliziny, quinolinyl, isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl,
25 quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, and phenoxazinyl;

Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halogen, hydroxyl, hydroxymethyl, nitro, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl, O-[(C₁-C₄)-straight or branched alkyl], O-benzyl, O-phenyl, 1,2-methylenedioxy, amino, carboxyl, N-[(C₁-C₅)-straight or branched alkyl or (C₂-C₅)-straight or branched alkenyl] carboxamides, N,N-di-[(C₁-C₅)-straight or branched alkyl or (C₂-C₅)-straight or branched alkenyl] carboxamides, N-morpholinecarboxamide, N-benzylcarboxamide, N-thiomorpholinocarboxamide, N-picolinoylcarboxamide, O-W, CH₂-(CH₂)_p-W, O-(CH₂)_p-W, (CH₂)_p-O-W, and

10 CH=CH-W;

W is 4-methoxyphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrazyl, quinolyl, 3,5-dimethylisoxazolyl, isoxazolyl, 2-methylthiazoyl, thiazoyl, 2-thienyl, 3-thienyl, or pyrimidyl; p is 0-2;

15


Q and A are independently hydrogen, Ar, (C₁-C₁₀)-straight or branched alkyl, (C₂-C₁₀)-straight or branched alkenyl or alkynyl, provided the carbon atom having the triple bond in the alkynyl group is not directly bonded to the nitrogen atom of the core; (C₅-C₇)-cycloalkyl substituted (C₁-C₆)-straight or branched alkyl,

20 (C₂-C₆)-straight or branched alkenyl or alkynyl, (C₅-C₇)-cycloalkenyl substituted (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, or Ar-substituted (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl wherein, in each case, any one of the CH₂ groups of said alkyl, alkenyl or alkynyl chains may be optionally replaced by a heteroatom selected from the group

25 consisting of O, S, SO, SO₂, N, and NR, wherein R is selected from the group consisting of hydrogen, (C₁-C₄)-straight or branched alkyl, (C₂-C₄)-straight or branched alkenyl or alkynyl, and (C₁-C₄)-bridging alkyl wherein a bridge is formed

8

between the nitrogen and a carbon atom of said heteroatom-containing chain to form a ring, and wherein said ring is optionally fused to an Ar group; or



5

G is hydrogen, (C₁-C₆)-straight or branched alkyl or (C₂-C₆)-straight or branched alkenyl or alkynyl; and

T is Ar or substituted 5-7 membered cycloalkyl with substituents at positions 3 and 4

10 which are independently selected from the group consisting of oxo, hydrogen, hydroxyl, O-(C₁-C₄)-alkyl, or O-(C₂-C₄)-alkenyl.

A preferred embodiment are compounds of formula I wherein

15 J and K are taken together to form a pyrrolidine ring;

the stereochemistry at carbon 1 is S;

B and C are each hydrogen;

X is F₂ or O;

Z is O or CH₂;

20 D is 3, 4, 5-trimethoxyphenyl or t-butyl;

A is 3-(3-pyridyl)propyl or 4-(3-pyridyl)butyl; and

Q is phenyl-substituted (C₁-C₆)-straight or branched chain alkyl, wherein phenyl is optionally substituted with one to three substituents independently selected from

C₁-C₆ alkyl, O-(C₁-C₆) alkyl, carboxyl and trifluoromethyl, wherein said alkyl is

25 straight or branched.

Another preferred embodiment are compounds of formula I wherein

Q is

- 3-phenylpropyl;
- 3-(3,4,5-trimethoxyphenyl)propyl;
- 5 2-(3,4-dimethoxyphenyl)ethyl;
- 2-phenylethyl;
- 4-phenylbutyl;
- 4-(3,4-dimethoxyphenyl)butyl;
- 3-(4-carboxyphenyl)propyl;
- 10 2-(3-methoxyphenyl)ethyl;
- 2-(3-trifluoromethylphenyl)ethyl, or
- 3-(4-methoxyphenyl)propyl.

- Another aspect of the present invention provides for a pharmaceutical
- 15 composition which comprises as an active ingredient an amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, effective for stimulating neurite growth in nerve cells, and one or more pharmaceutically acceptable carriers, excipients or diluents thereof.


- 20 Another aspect of the present invention provides for a method for stimulating neurite growth in nerve cells comprising the step of contacting said nerve cells with a composition comprising a neurotrophic amount of a compound of formula I with affinity for an FK-506 binding protein.

- 25 Another aspect of the present invention provides for a method for stimulating neurite growth in nerve cells comprising the step of contacting said nerve cells with a composition comprising a neurotrophic amount of a compound of formula I with affinity for FKBP12.

GENERAL SUMMARY OF COMPOUND PREPARATION


The carbamates of this invention are best prepared according to the general scheme shown below. Aminoalcohol nitrogens are protected by acylation with di-*t*-butyl carbonate and the resulting carbamate alcohols are treated with *p*-nitrophenyl chloroformate to give activated carbonates. These can be coupled with amines, in the presence of catalytic hydroxybenzotriazole, to give Boc-protected carbamates. The Boc protecting group is removed by exposure to hydrogen chloride, and the resulting amine is acylated to give the target compounds.

10




Alternatively, the carbamates of the present invention can also be prepared according to the general scheme shown below. Aminoalcohols can be selectively acylated at the amino group with an activated carboxyl derivative. Treatment with *p*-nitrophenyl chloroformate, and coupling with an amine is carried out as described above to give the target compounds.

15




Intermediates used to prepare secondary and tertiary alkyl-substituted carbamates can
 5 be prepared by Grignard additions to aldehydes and esters as shown below.




For differently-substituted compounds (i.e., where B and C are different), the
 10 mono-alkylated intermediate can be re-oxidized and treated with a second Grignard reagent.

12




The secondary amine intermediates utilized in this invention can be prepared either by reductive amination of an aldehyde with a primary amine as shown below:

5




or by acylation of an amine, followed by reduction of the amide with diborane or lithium aluminum hydride:

10




Diaryl secondary amine intermediates can be prepared by transition metal mediated coupling reactions such as those shown below.

15



OR




where QX is appropriately activated by electron withdrawing substituents.


20

Enamine intermediates may be prepared by condensation of primary amines with aldehydes and ketones as shown below.


13



5 The 2,2-difluoroacetic acids are synthesized by fluorination of the parent keto compound with diethylaminosulfur trifluoride. The N,N-diethylamide is obtained as the major product, but it is easily converted to the desired acid by alkaline hydrolysis.




10 The 2,2-difluoroacetic acids are converted to the corresponding acid chlorides using oxalyl chloride and catalytic dimethylformamide in methylene chloride.



15 The diamides of this invention are best prepared according to the general scheme shown below. N-Boc-proline methyl ester is reduced to the corresponding aldehyde and coupled with the appropriate Wittig reagent to give the methyl acrylate which is then reduced to the 3-pyrrolidinylpropionate. The methyl ester is hydrolyzed to the acid, which is coupled to the secondary amines described above. Removal of the


20 Boc protecting group and coupling to the keto- or difluoroacids are carried out as described above for the carbamate examples.

14



Alternatively, the aldehydes in the scheme shown above can be prepared by oxidation of the corresponding alcohols.

5



PREPARATION OF INTERMEDIATES

10 N-t-Boc-L-pyrrolidinylmethyl-p-nitrophenylcarbonate

A stirred solution of L-prolinol (10.21 g, 100 mmoles) in methylene chloride (75 mL) at room temperature (rt) was treated with t-butyl pyrocarbonate (22.03 g, 1 equiv.) at such a rate as to cause only a gentle reflux. Following addition, the mixture was

stirred at rt for 1.5 h and then the solvent was removed by evaporation. The residue was dried *in vacuo* for several hours and then dissolved in dry methylene chloride (150 mL). The solution was cooled to 0°C and treated with p-nitrophenylchloroformate (20.37 g, 1 equiv.) and pyridine (8.2 mL, 1 equiv.). After stirring at rt overnight the solvent was removed by evaporation and the residue dissolved as far as possible in ether (250 mL). The solid pyridine hydrochloride was removed by filtration and the filtrate washed with water, 2% phosphoric acid, water, and brine, dried over sodium sulfate and evaporated to give a thick, pale-yellow oil (27.90 g, 75%) that was pure enough by NMR to carry on without further purification.

¹H-NMR δ (CDCl₃) 1.46 (9H, s), 1.99 (4H, m), 3.41 (2H, m), 4.07 and 4.22 (1H, 2 x m), 4.31 (2H, d), 7.39 (2H, d), 8.25 (2H, d).

Mass Spec. (ESI): 367.4 (MH)⁺.

N-t-Boc-(±)-piperidinylmethyl-p-nitrophenylcarbonate

This was prepared as described in the prior preparation, from racemic 2-hydroxymethylpiperidine giving 23.62 g of the product (95%) as a tan solid.

¹H-NMR δ (CDCl₃) 1.43 (11H, m and s), 1.68 (4H, m), 2.83 (1H, t), 4.02 (1H, brd), 4.31 (1H, ABq), 4.44 (1H, t), 4.66 (1H, m), 7.40 (2H, d), 8.24 (2H, d).

Mass Spec. (ESI): 381.3 (MH)⁺.

N-Trimethylpyruvyl-L-pyrrolidinylmethanol

A stirred solution of trimethylpyruvic acid (1.37 g, 10.50 mmol), N-hydroxybenzotriazole (HOBt) (1.61 g, 1 equiv.) and L-prolinol (1.24 mL, 1.2 equiv.) in tetrahydrofuran (100 mL) at 0°C was treated with 0.5M dicyclohexylcarbodiimide in methylene chloride (25.2 mL, 1.2 equiv.). The mixture was stirred at rt overnight and then the solid urea by-product was removed by filtration. The filtrate was concentrated *in vacuo* and the residue dissolved in 1:1 ether/ethyl acetate. The solution was washed

with water and brine, dried over sodium sulfate and evaporated to give a white, waxy solid (1.48 g, 66%).

$^1\text{H-NMR}$ δ (CDCl_3) 1.03 (9H, s), 1.72 (1H, m), 1.91 (2H, m), 2.04 (1H, m), 3.45 (1H, ABq), 3.51 (1H, ABq), 3.63 (1H, ABq), 3.76 (1H, m), 4.02 (1H, ABq).

5 Mass Spec. (ESI): 214.3 (MH)⁺.

N-Trimethylpyruvyl-L-pyrrolidinylmethyl-p-nitrophenylcarbonate

A stirred solution of the alcohol from Example 3 (242.3 mg, 1.136 mmoles) in methylene chloride (10 mL) at 0°C was treated with p-nitrophenyl chloroformate
10 (274.8 mg, 1.2 equiv.) and pyridine (0.11 mL, 1.2 equiv.). The mixture was stirred at rt overnight. Thin layer chromatography indicated ca. 30% unreacted starting material. More p-nitrophenyl chloroformate (114.5 mg, 0.5 equiv.) and pyridine (0.046 mL, 0.5 equiv.) were added. Stirring was continued overnight at rt. The mixture was then concentrated *in vacuo* and the residue dissolved in 1:1 ether/ethyl
15 acetate. The solution was washed with 2% phosphoric acid, water and brine, dried over sodium sulfate and evaporated. The residue was chromatographed on silica gel, eluting with 35% ethyl acetate/hexane, to give the product as a white, waxy solid (288 mg, 67%).

$^1\text{H-NMR}$ δ (CDCl_3) 1.28 (9H, d), 1.95 (2H, m), 2.04 (1H, m), 2.12 (1H, m), 3.42
20 (2H, m), 4.33 (1H, m), 4.51 (2H, m), 7.40 (2H, d), 8.29 (2H, d).

Mass Spec. (ESI): 379.3 (MH)⁺.

3-(3-Pyridyl)propylammonium p-toluenesulfonate

A stirred solution of 3-(3-pyridyl)propanol (3.5 mL, 27.12 mmoles) in methylene
25 chloride (15 mL) at 0°C was treated with p-toluenesulfonyl chloride (6.7219 g, 1.3 equiv.) and diisopropylethylamine (6.14 mL, 1.3 equiv.). After 10 minutes, the mixture was warmed to room temperature. After a further 4 hours, the solvent was removed under reduced pressure and the residue partitioned between ethyl acetate and

water. The organic phase was washed with brine, dried over sodium sulfate and evaporated. the residue was chromatographed on silica, eluting with ethyl acetate, to give a thick, pale-yellow oil which was immediately dissolved in saturated ammonia/methanol (165 mL). This was left to stand at room temperature for 5 days at
5 which time the solvent was removed under reduced pressure. The residue was dried under high vacuum and then dissolved as far as possible in ether (150 mL). The resulting solid product was collected by filtration, washed with ether, and dried in vacuo (6.3051 g, 76%).

¹H-NMR δ (DMSO-d₆) 1.84 (m, 2H), 2.28 (s, 3H), 2.64 (t, 2H), 2.80 (t, 2H), 7.11
10 (d, 2H), 7.35 (ABq, 1H), 7.49 (d, 2H), 7.63 (m, 1H), 7.69 (br, 3H), 8.41 (m, 2H).

3-(3-Pyridyl)propyl-3-phenylpropylamine

A stirred mixture of 3-(3-pyridyl)propionaldehyde (4.40 g, 32.5 mmoles) and 3-phenylpropylamine (5.30 g, 1.2 equiv.) in methanol at rt was treated with sodium
15 cyanoborohydride (1 g, 0.5 equiv.). After 1 h at rt the mixture was further treated with solid potassium hydroxide (3.6 g, 2 equiv.) and stirring was continued for 2 h. The resulting mixture was partitioned between brine and ether. The aqueous layer was extracted with ether and the combined ether extracts were extracted with 6N HCl (3x). These aqueous extracts were basified with solid potassium hydroxide and extracted
20 with ether (3x), dried over sodium sulfate and concentrated *in vacuo*. The residue was distilled under high vacuum to give the product as a colorless oil (2.44 g, 30%), b.p. ca. 162°C/0.5 mm Hg.

¹H-NMR δ (CDCl₃) 1.77 (4H, m), 2.42 (1H, m), 2.63 (8H, m), 7.18 (4H, m), 7.26
(2H, m), 7.49 (1H, t), 8.43 (2H, m).

25

The following compounds were prepared essentially as described for the previous preparation:

3-(3-Pyridyl)propyl-4-phenylbutylamine

¹H-NMR δ (CDCl₃) 8.39 (br s, 1 H); 7.44 (d, 1 H); 7.29-7.14 (m, 7 H); 2.89-2.77 (m, 4 H); 2.66-2.57 (m, 4 H); 2.17 (br s, 1 H); 1.75 (q, 2 H).

5 3-(3-Pyridyl)propyl-2-phenylethylamine

¹H-NMR δ (CDCl₃) 8.41 (m, 1 H); 7.45 (d, 1 H); 7.31-7.13 (m, 7 H); 2.95-2.71 (m, 2 H); 2.68-2.54 (m, 4 H); 2.47 (t, 2 H); 2.05 (m, 1 H); 1.67-1.41 (m, 4 H).

3-(3-Pyridyl)propyl-benzylamine

10 ¹H-NMR δ 8.44 (m, 1 H); 7.44 (d, 1 H); 7.35-7.19 (m, 7 H); 3.77 (s, 2 H); 2.68 (m, 4 H); 1.82 (q, 2 H); 1.78 (br s, 1 H).

4-(3-Pyridyl)butyl-4-phenylbutylamine

15 ¹H-NMR δ 8.41 (m, 0.5 H); 7.24 (m, 1.5 H); 7.34-7.07 (m, 7 H); 2.80 (m, 2 H); 2.66 (m, 8 H); 1.88 (m, 2 H); 1.78-1.42 (m, 5 H).

4-(3-Pyridyl)butyl-3-phenylpropylamine

¹H-NMR δ 8.41 (m, 1 H); 7.23 (m, 1 H); 7.34-7.07 (m, 7 H); 2.80 (m, 2 H); 2.61 (t, 4 H); 2.53 (t, 4 H); 1.83-1.47 (br m, 7 H).

20

4-(3-Pyridyl)butyl-2-phenylethylamine

¹H-NMR δ 8.41 (m, 1 H); 7.45 (d, 1 H); 7.31-7.13 (m, 7 H); 2.95-2.71 (m, 2 H); 2.68-2.54 (m, 4 H); 2.47 (t, 2 H); 2.05 (m, 1 H); 1.67-1.41 (m, 4 H).

25 4-(3-Pyridyl)butyl-benzylamine

¹H-NMR δ 8.43 (m, 1.5 H); 7.42 (m, 0.5 H); 7.38-7.17 (m, 7 H); 3.49 (s, 2 H); 2.53 (q, 2 H); 2.39 (q, 2 H); 1.61-1.41 (m, 5 H).

2-(3-Indolyl)ethyl-2-phenylethylamine

$^1\text{H-NMR}$ δ 7.97 (bs, 1 H); 7.64 (d, J = 8.2 Hz, 1 H); 7.38 (d, J = 8.3 Hz, 1 H); 7.29-7.07 (m, 7 H); 6.95 (d, J = 2.4 Hz, 1 H); 2.99 (s, 4 H); 2.92 (t, J = 7.0 Hz, 2 H); 2.80 (t, J = 6.8 Hz, 2 H).

5 Mass Spec. (ESI): 265.2 (MH)⁺.

(E) Ethyl 4-carboxycinnamate

4-Carboxybenzaldehyde (1.0 g) was added to sodium hydride (170 mg) in 100 mL tetrahydrofuran at 0°C. In a separate flask, triethyl phosphonoacetate (1.60 mL) was
10 added to sodium hydride (200 mg) in 100 mL tetrahydrofuran at 0°C. The phosphonate anion solution was cannulated into the slurry of the carboxylate anion and the resulting thick white slurry was stirred for 16 hours. Water (50 mL) was added followed by concentrated hydrochloric acid until the pH was 1. The precipitate was collected by filtration, dissolved in ether and dried with magnesium sulfate.
15 Evaporation under reduced pressure gave the product (1.47 g; 100%) as a white solid.
 $^1\text{H-NMR}$ δ (CDCl₃) 8.13 (d, J = 8.2 Hz, 2 H); 7.73 (d, J = 16.3 Hz, 1 H); 7.64 (d, J = 8.0 Hz, 2 H); 6.52 (d, J = 16.1 Hz, 1 H); 4.29 (q, J = 7.3 Hz, 2 H); 1.35 (t, J = 7.5 Hz, 3 H).

20 Ethyl 3-(4-carboxyphenyl) propionate

A solution of (E) ethyl 4-carboxycinnamate in ethanol was treated with 20% palladium hydroxide on carbon (500 mg). The mixture was stirred under a hydrogen atmosphere for 2 days and was then filtered through celite to remove inorganics. Evaporation gave ethyl 3-(4-carboxyphenyl) propionate as a white solid (1.19 g; 80% yield).
25 $^1\text{H-NMR}$ δ (CDCl₃) 8.04 (d, J = 8.2 Hz, 2 H); 7.31 (d, J = 8.1 Hz, 2 H); 4.13 (q, J = 7.1 Hz, 2 H); 3.04 (t, J = 7.7 Hz, 2 H); 2.66 (t, J = 7.7 Hz, 2 H); 1.23 (t, J = 7.3 Hz, 3 H).

Benzyl 4-(3-hydroxypropyl)benzoate

A stirred solution of ethyl 3-(4-carboxyphenyl) propionate in 200 mL tetrahydrofuran was treated with lithium aluminum hydride (400 mg). After 4 hours, excess hydrochloric acid was added and the mixture was extracted with ether. The organic layer was dried with magnesium sulfate and the solvent was removed *in vacuo*. This residue was dissolved in 2 mL dimethylformamide and to this was added benzyl bromide (700 μ L) and potassium carbonate (1.0 g). The reaction was stirred for 16 hours and then diluted with water (20 mL). The solution was saturated with sodium chloride and extracted with ethyl acetate. The organic layer was dried with magnesium sulfate and purified by silica gel chromatography to give the product as a clear oil (385 mg, 27%).

$^1\text{H-NMR}$ δ (CDCl_3) 8.00 (d, $J = 8.6$ Hz, 2 H); 7.48-7.30 (m, 5 H); 7.27 (d, $J = 8.6$ Hz, 2 H); 5.35 (s, 2 H); 3.68 (t, $J = 7.1$ Hz, 2 H); 2.77 (t, $J = 7.4$ Hz, 2 H); 1.90 (m, 2 H). The product was oxidized by either Dess-Martin periodinane or Swern methods to give the corresponding aldehyde which could be used in reductive aminations according to the procedure described for 3-(3-Pyridyl)-propyl-3-phenylpropylamine. The amine resulting from reductive amination with 3-(3-pyridyl)propyl amine was not fully characterized but showed Mass Spec. (ESI): 389.3 (MH) $^+$.

20

3-(3-Pyridyl)propyl-4-(3,4-dimethoxyphenyl)butylamine

A stirred solution of 3-(3-pyridyl)propylamine (517.8 mg, 1.679 mmoles), 4-(3,4-dimethoxyphenyl)butanoic acid (376.5 mg, 1 equiv.) and N-hydroxysuccinimide (193.2 mg, 1.1 equiv.) in methylene chloride (15 mL) at 0°C was treated with EDC hydrochloride (353.1 mg, 1.1 equiv.). The mixture was allowed to warm to room temperature after 10 minutes. Stirring was continued for 16 hours and then the solvent was removed under reduced pressure. The residue was partitioned between ethyl acetate and water. The aqueous phase was re-extracted with ethyl acetate. The

combined organic phases were washed with water and brine, dried over sodium sulfate and evaporated to give a colorless oil. This crude amide was dissolved in dry tetrahydrofuran (5 mL) under argon. The solution was cooled to 0°C and then treated with 1 M borane-tetrahydrofuran complex (7.0 mL, 4 equiv.). The mixture was heated at reflux for 16 hours and then cooled to room temperature. Dry methanol (10 mL) was added carefully to the stirred mixture to avoid foaming. The resulting solution was evaporated under reduced pressure and the residue dissolved in dry methanol (20 mL). The mixture was heated at reflux for 4 hours and then evaporated. The crude amine was carried on without purification.

¹H-NMR δ (CDCl₃) 1.61 (m, 6H), 2.12 (br, 1H), 2.60 (m, 8H), 3.85 (s, 3H), 3.86 (s, 3H), 6.71 (m, 3H), 7.18 (ABq, 1H), 7.47 (d, 1H), 8.41 (m, 2H).

The following compounds were prepared essentially as described for the previous preparation:

15

3-(3-Pyridyl)propyl-3-(3,4,5-trimethoxyphenyl)propylamine

¹H-NMR δ (CDCl₃) 1.80 (m, 4H), 2.63 (m, 8H), 3.80 (s, 3H), 3.82 (2 x s, 6H), 6.39 (d, 2H), 7.18 (ABq, 1H), 7.47 (m, 1H), 8.40 (m, 2H).

20 **3-(3-Pyridyl)propyl-3-(3-methoxyphenyl)propylamine**

¹H-NMR δ (CDCl₃) 1.83 (m, 4H), 2.64 (m, 8H), 3.75 (s, 3H), 6.75 (m, 3H), 7.17 (m, 2H), 7.44 (m, 1H), 8.37 (m, 2H).

3-(3-Pyridyl)propyl-3-(4-methoxyphenyl)propylamine

25 ¹H-NMR δ (CDCl₃) 1.78 (m, 4H), 2.57 (m, 8H), 3.72 (s, 3H), 6.77 (ABq, 2H), 7.06 (ABq, 2H), 7.16 (ABq, 1H), 7.42 (m, 1H), 8.39 (m, 2H).

3,4,5-Trimethoxyphenyl-2-oxoacetyl chloride

A stirred suspension of 3',4',5'-trimethoxyphenyl-2-oxoacetic acid (1.60 g, 6.66 mmol) in dry methylene chloride (26 mL) at rt was treated with 2M oxalyl chloride in methylene chloride (14 mL, 4 equiv.) and dry dimethylformamide (1 drop). After 3 h the solvents were evaporated. The residue was flushed with dry methylene chloride (3 x 50 mL) and dried *in vacuo* for 2 h during which time a solid formed. The crude acid chloride was carried on without further purification.

2,2-Difluoro-(3,4,5-trimethoxyphenyl)acetic acid.

- 10 To a stirred solution of 3,4,5-trimethoxyphenyl-2-oxoacetic acid (3.81 g, 15.8 mmol) in anhydrous methylene chloride (30 mL) at room temperature was added diethylaminosulfurtrifluoride (DAST) (20.4 g, 127 mmol) under nitrogen and the mixture was stirred overnight. The mixture was then cooled in an ice bath and excess DAST was quenched by dropwise addition of water. Ethyl acetate (300 mL) was added and the organic layer was washed with saturated aqueous sodium bicarbonate (2 x 100 mL) followed by water (100 mL). The residue obtained after drying and evaporation was purified by silica gel chromatography, eluting with hexane/ethyl acetate (9:1 to 7:3), to give the N,N-diethylamide derivative (2.10 g, 6.62 mmol, 42%) as a pale yellow solid. ¹H-NMR: 6.77 (s, 2H), 3.88 (s, 9H), 3.45 (q, *J* = 7.0 Hz, 2H), 3.25 (q, *J* = 7.0 Hz, 2H), 1.20 (t, *J* = 7.0 Hz, 3H), 1.10 (t, *J* = 6.9 Hz, 3H). Bicarbonate washing after acidification and extractive work up with ethyl acetate gave the crude title compound. Purification by reversed phase column (C18) chromatography eluting with water/methanol/trifluoroacetic acid (69.9:30:0.1) gave the pure 2,2-difluoro acid (0.616 g, 2.34 mmol, 15%) as a white solid. ¹H-NMR: 6.85 (s, 2H), 3.90 (s, 6H), 3.89 (s, 3H). Anal. C: 50.59, H: 4.72, F: 14.24 (found), C: 50.39, H: 4.61, F: 14.49 (calcd). The N,N-diethylamide (2.00 g, 6.30 mmol) obtained above was hydrolyzed to the title acid by heating a solution in ethanol (5 mL) with 10% sodium hydroxide (13 mL) at reflux for 4 h. Acidification followed by extractive work
- 25

up with ethyl acetate gave the crude acid which was purified as described above to give 1.51 g of the title compound as a white solid.

2,2-Difluoro-3,4,5-trimethoxyphenylacetyl chloride

- 5 This was prepared as described above for 3,4,5-trimethoxyphenyl-2-oxoacetyl chloride from the corresponding carboxylic acid and was used without chromatographic purification.

N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)

10 **propyl-3-phenylpropylamino]carbamate**

- A stirred solution of 3-(3-pyridyl)propyl-3-phenylpropylamine (342.5 mg, 1.346 mmol) and N-Boc-L-prolinyl-p-nitrophenylcarbonate (612.2 mg, 1.24 equiv.) in methylene chloride (7 mL) at rt was treated with N-hydroxybenzotriazole (37 mg, 0.2 equiv.) and diisopropylethylamine (0.235 mL, 1 equiv.). After stirring overnight at rt, 15 the mixture was diluted with ethyl acetate and the solution was washed with 1M sodium hydroxide (2x), water and brine, dried over sodium sulfate, and concentrated *in vacuo*. The residue was chromatographed on silica gel, eluting with ethyl acetate, to give the product as a pale-yellow oil (500.6 mg, 77%).

- ¹H-NMR δ (CDCl₃) 1.42 (9H, s), 1.83 (8H, m), 2.58 (4H, brt), 3.27 (6H, m), 4.04 20 (3H, m), 7.17 (4H, m), 7.25 (2H, m), 7.50 (1H, m), 8.42 (2H, brs). Mass Spec. (ESI): 482.6 (MH)⁺.

The following compounds were prepared essentially as described for the previous preparation:

N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-N'-3-pyridylmethylcarbamate

¹H-NMR δ (CDCl₃) 1.43 (9H, s), 1.82 (4H, m), 3.33 (2H, brs), 4.10 (3H, m), 4.38 (2H, d), 5.22 (1H, br), 7.24 (1H, ABq), 7.60 (1H, d), 8.48 (2H, brs).

5

N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-N'-2-phenylethylcarbamate

¹H-NMR δ (CDCl₃) 1.42 (9H, s), 1.81 (4H, m), 2.78 (2H, t), 3.28 (2H, m), 3.40 (2H, m), 3.95 (2H, m), 4.10 (1H, m), 4.77 (1H, br), 7.21 (5H, m).

10 **N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-benzylamino]carbamate**

¹H-NMR δ (CDCl₃) 8.41 (d, 0.5 H); 8.38 (s, 0.5 H); 7.43 (m, 1 H); 7.21 (m, 7H); 4.17 (m, 2 H); 3.91 (m, 1 H); 3.33 (br s, 2 H); 3.20 (br s, 2 H); 2.55 (m, 2 H); 1.80 (br m, 6 H); 1.40 (s, 9 H).

15

N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-2-phenylethylamino]carbamate

¹H-NMR δ (CDCl₃) 8.41 (m, 1 H); 7.43 (m, 1 H); 7.21 (m, 7 H); 4.17 (m, 2 H); 3.91 (m, 1 H); 3.30 (br m, 6 H); 2.81 (d, 2 H); 2.57 (d, 2 H); 1.81 (m, 6 H); 1.42 (s, 9 H).

20

N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-4-phenylbutylamino]carbamate

¹H-NMR δ (CDCl₃) 8.41 (m, 1 H); 7.43 (m, 1 H); 7.20 (m, 7 H); 4.50 (m, 1 H); 4.15 (m, 3 H); 3.41 (m, 2 H); 3.19 (m, 2 H); 2.81 (m, 3 H); 2.59 (m, 2 H); 1.79 (m, 2 H); 1.62 (m, 10 H); 1.41 (s, 9 H).

25

**N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)
butyl-2-phenylethylamino]carbamate**

¹H-NMR δ (CDCl₃) 8.42 (m, 1 H); 7.43 (m, 1 H); 7.20 (m, 7 H); 4.20 (m, 2 H);
3.87 (m, 1 H); 3.26-2.95 (m, 6 H); 2.79-2.43 (br m, 4 H); 1.84-1.63 (br m, 8 H);
5 1.41 (s, 9 H).

**N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)
butyl-4-phenylbutylamino]carbamate**

¹H-NMR δ (CDCl₃) 1.41 (9H, s), 1.57 (6H, m), 1.81 (4H, m), 2.60 (4H, brt), 3.21
10 (6H, m), 3.90 (1H, m), 3.97 (2H, m), 4.13 (1H, m), 7.19 (6H, m), 7.47 (1H, d),
8.41 (2H, brs).

**N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)
(propyl-3-(m-methoxyphenyl)propylamino]carbamate**

15 ¹H-NMR δ (CDCl₃) 1.42 (s, 9H), 1.81 (m, 8H), 2.58 (m, 4H), 3.23 (m, 6H), 3.79
(s, 3H), 4.04 (m, 3H), 6.75 (m, 3H), 7.20 (m, 2H), 7.49 (brt, 1H), 8.42 (brs, 2H).
Mass Spec. (ESI): 512.42 (MH)⁺.

**N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)
propyl-3-(p-methoxyphenyl)propylamino]carbamate**

20 ¹H-NMR δ (CDCl₃) 1.45 (s, 9H), 1.82 (m, 8H), 2.57 (m, 4H), 3.23 (m, 6H), 3.77
(s, 3H), 4.03 (m, 3H), 6.81 (d, 2H), 7.06 (brd, 2H), 7.21 (ABq, 1H), 7.48 (brt,
1H), 8.42 (brs, 2H).
Mass Spec. (ESI): 512.41 (MH)⁺.

**N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)
propyl-3-(3,4,5-trimethoxyphenyl)propylamino]carbamate**

¹H-NMR δ (CDCl₃) 1.43 (s, 9H), 1.82 (m, 8H), 2.55 (m, 4H), 3.28 (m, 6H), 3.79 (s, 3H), 3.81 (s, 3H), 4.02 (m, 3H), 6.36 (brs, 2H), 7.19 (ABq, 1H), 7.44 (m, 1H),
5 8.41 (brs, 2H).

Mass Spec. (ESI): 572.46 (MH)⁺.

**N-t-Butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)
propyl-4-(3,4-dimethoxyphenyl)butylamino]carbamate**

10 ¹H-NMR δ (CDCl₃) 1.43 (s, 9H), 1.53 (m, 4H), 1.81 (m, 6H), 2.54 (m, 4H), 3.22 (m, 6H), 3.82 (s, 3H), 3.84 (s, 3H), 4.07 (m, 3H), 6.66 (brs, 2H), 6.75 (m, 1H),
7.19 (ABq, 1H), 7.46 (m, 1H), 8.41 (brs, 2H).

Mass Spec. (ESI): 556.44 (MH)⁺.

15 **N-t-Butyloxycarbonyl-(±)-piperidinylmethyl-[N',N'-3-(3-pyridyl)
propyl-3-phenylpropylamino]carbamate**

¹H-NMR δ (CDCl₃) 1.42 (11H, m and s), 1.54 (4H, m), 1.82 (4H, m), 2.56 (4H, m), 2.81 (1H, q), 3.22 (4H, m), 3.97 (2H, brd), 4.12 (1H, ABq), 4.18 (1H, m),
4.44 (1H, br), 7.17 (4H, m), 7.22 (2H, t), 7.45 (1H, brd), 8.42 (2H, brs).

20

**N-t-Butyloxycarbonyl-(±)-piperidinylmethyl-[N',N'-3-(3-pyridyl)
propyl-benzylamino]carbamate**

¹H-NMR δ (CDCl₃) 8.41 (d, 0.5 H); 8.38 (s, 0.5 H); 7.42 (br t, 1 H); 7.21 (m, 7 H); 4.42 (m, 3 H); 4.23 (m, 1 H); 4.17 (m, 4 H); 3.22 (m, 2 H); 2.81 (m, 1 H);
25 2.50 (m, 2 H); 1.80 (m, 2 H); 1.55 (br s, 4 H); 1.41 (s, 9 H).

**N-t-Butyloxycarbonyl-(±)-piperidinylmethyl-[N',N'-3-(3-pyridyl)
propyl-2-phenylethylamino]carbamate**

¹H-NMR δ (CDCl₃) 8.41 (s, 1 H); 7.43 (br t, 1 H); 7.21 (m, 7 H); 4.12 (m, 2 H);
3.27 (m, 5 H); 2.60 (m, 4H); 1.81 (br s, 6 H); 1.57 (br s, 4 H); 1.42 (s, 9 H).

5

**N-t-Butyloxycarbonyl-(±)-piperidinylmethyl-[N',N'-3-(3-pyridyl)
propyl-4-phenylbutylamino]carbamate**

¹H-NMR δ (CDCl₃) 8.41 (s, 1 H); 7.42 (m, 1 H); 7.20 (m, 7 H); 4.42 (m, 1 H);
4.15 (m, 2 H); 4.01 (m, 1 H); 3.21 (m, 4 H); 2.82 (br t, 1 H); 2.68 (br s, 4 H);
1.81 (m, 2 H); 1.55 (m, 10 H); 1.41 (s, 9 H).

10

**(S)-2-(2-Methoxycarbonylvinyl)-piperidine-1-carboxylic acid t-butyl
ester**

A stirred solution of (S)-N-(t-butoxycarbonyl)-pipecolic acid methyl ester (2.04 g,
15 8.39 mmol) in methylene chloride (75 mL) at -78°C under nitrogen was treated with
diisobutylaluminum hydride (1.0 M in toluene, 14.2mL, 14.2 mmol). After 1.25 h a
second aliquot of diisobutylaluminum hydride (2.00 mL, 2.00 mmol) was added. The
mixture was stirred for an additional 30 min. and then quenched by the addition of a
20% aqueous Rochelle's salt solution (100 mL) followed by vigorous stirring for 1 h.
20 The aqueous layer was extracted with methylene chloride and the combined organic
extracts were dried over magnesium sulfate, filtered, and concentrated. The crude
aldehyde was dissolved in methylene chloride (100 mL) and treated with methyl
(triphenylphosphoranylidene)acetate (4.00 g, 12.0 mmol). After 15 hours (h) the
reaction mixture was concentrated and purified by flash chromatography to give the
25 desired product (1.46 g, 65%).

¹H-NMR δ (CDCl₃) 1.35 (m, 2 H), 1.45 (s, 9 H), 1.62 (m, 2 H), 1.78 (m, 2 H),
2.80 (t, 1 H, *J* = 11.6), 3.74 (s, 3 H), 3.98 (d, 1 H, *J* = 13.1), 4.94 (s, 1 H), 5.81
(dd, 1 H, *J* = 1.7, 15.9), 6.88 (dd, 1 H, *J* = 4.0, 15.9).

(S)-2-(2-Methoxycarbonylvinyl)-pyrrolidine-1-carboxylic acid t-butyl ester

To a solution of oxalyl chloride (2.0M in methylene chloride, 9.60 mL, 19.2 mmol) in methylene chloride (100 mL) at -60°C was added a solution of dimethyl sulfoxide (3.01 g, 38.6 mmol) in methylene chloride (10 mL) and the mixture was stirred for 5 min. A solution of (S)-(-)-1-(t-butoxycarbonyl)-2-pyrrolidinemethanol (3.08 g, 15.3 mmol) in methylene chloride (20 mL) was added dropwise over 5 min. After 30 min, triethylamine (12.8 mL, 91.8 mmol) was added and the reaction mixture was stirred at -60°C for 15 min and then warmed to rt for a further 10 min. Water (50 mL) was added and the mixture extracted with methylene chloride. The combined organic extracts were dried over magnesium sulfate, filtered, and concentrated. The crude residue was dissolved in methylene chloride (100 mL) and treated with methyl (triphenylphosphoranylidene)acetate (1.4 eq). After 15 h the reaction mixture was concentrated and purified by silica gel chromatography to give the desired product (3.54 g, 90%).

¹H-NMR δ (CDCl₃) 1.41 (s, 9 H), 1.85 (m, 3 H), 2.06 (m, 1 H), 3.42 (m, 2 H), 3.73 (s, 3 H), 4.42 (m, 1 H), 5.82 (d, 1 H, J = 15.4), 6.82 (m, 1 H).

(S)-2-(2-Methoxycarbonyl-ethyl)-piperidine-1-carboxylic acid t-butyl ester

To a solution of (S)-2-(2-methoxycarbonylvinyl)-piperidine-1-carboxylic acid t-butyl ester (1.46 g, 5.41 mmol) in 1:1 methanol:methylene chloride (40 mL) was added 10% palladized charcoal (0.129 g). The mixture was hydrogenated (59 psi) for 5.5 h. The catalyst was removed by filtration through a bed of celite and the solution concentrated *in vacuo*. Purification by silica gel chromatography provided the product (1.10 g, 75%).

¹H-NMR δ (CDCl₃) 1.44 (s, 9 H), 1.60 (m, 7 H), 2.11 (m, 1 H), 2.30 (m, 2 H), 2.73 (t, 1 H, J = 13.5), 3.66 (s 3 H), 4.11 (m, 1 H), 4.24 (m, 1 H).

The following compound was prepared essentially as described for the previous preparation:

(S)-2-(2-Methoxycarbonyl-ethyl)-pyrrolidine-1-carboxylic acid t-butyl

5 **ester**

¹H-NMR δ (CDCl₃) 1.45 (s, 9 H), 1.65 (m, 2 H), 1.90 (m, 4 H), 2.32 (t, 2 H, $J = 7.9$), 3.32 (m, 2 H), 3.66 (s, 3 H), 3.80 (m, 1 H).

10 **N,N-(3-Phenylpropyl)-3-(3-pyridyl)propyl-2-[(S)-N'-t-butoxycarbonyl-2-piperidinyl]propionamide**

A solution of (S)-2-(2-methoxycarbonyl-ethyl)-piperidine-1-carboxylic acid t-butyl ester (0.544 g, 2.00 mmol) in tetrahydrofuran (15 mL) was treated with LiOH (1M, 15 mL). After 2.5 h the mixture was acidified with 1N HCl (pH 2) and extracted with methylene chloride. The combined organic extracts were dried over magnesium sulfate, filtered and concentrated. The crude product was dissolved in methylene chloride (10 mL) and cooled to 0°C. 3-(3-Pyridyl)propyl-3-phenylpropylamine (0.506 g, 1.99 mmol) in methylene chloride (5 mL) was added, followed by 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide (EDC) (0.414 g, 2.16 mmol), 1-hydroxybenzotriazole (0.292 g, 2.16 mmol), and triethylamine (0.560 mL, 4.02 mmol). The solution was allowed to warm to rt and stirred for 15 h. The reaction mixture was diluted with methylene chloride (100 mL). The organic layer was washed with 1N HCl, saturated sodium bicarbonate, and brine, then dried (MgSO₄), filtered, and concentrated *in vacuo*. Purification by flash chromatography yielded the product (0.755 g, 76%).

25 ¹H-NMR δ (CDCl₃) 1.43 (s, 9 H), 1.56 (m, 4 H), 1.65 (m, 2 H), 1.85 (m, 5 H), 2.15 (m, 3 H), 2.60 (m, 4 H), 2.72 (m, 1 H), 3.22 (m, 2 H), 3.33 (m, 2 H), 3.95 (m, 1 H), 4.23 (m, 1 H), 7.22 (m, 5 H), 7.53 (dd, 1 H, $J = 5.3, 7.8$), 7.97 (d, 1 H, $J = 7.8$), 8.50 (m, 2 H).

Mass Spec. (ESI): 494 (MH)⁺.

The following compound was prepared essentially as described for the previous preparation:

5


N,N-(3-Phenylpropyl)-3-(3-pyridyl)propyl-2-[(S)-N'-t-butoxycarbonyl-2-pyrrolidinyl]propionamide

¹H-NMR δ (CDCl₃) 1.43 (s, 9 H), 1.65 (m, 2 H), 1.83 (m, 8 H), 2.20 (m, 2 H),
2.62 (m, 3 H), 2.72 (m, 1 H), 3.27 (m, 6 H), 3.82 (m, 1 H), 7.25 (m, 5 H), 7.75 (m,
10 1 H), 8.02 (m, 1 H), 8.52 (m, 2 H).

Mass Spec. (ESI): 480 (MH)⁺.

EXAMPLES

15 **Example 1 N-Trimethylpyruvyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-phenylpropylamino]carbamate**




20 A mixture of 3-(3-pyridyl)-propyl-3-phenylpropylamine (0.38 g, 1.10 mmoles) and N-trimethylpyruvyl-L-prolinyl-p-nitrophenylcarbonate (0.46 g, 1.1 equiv.) in methylene chloride (2 mL) was stirred at rt over 4 days. The mixture was then washed with water, dried over magnesium sulfate and concentrated *in vacuo*. The residue was

chromatographed on silica gel, eluting with ethyl acetate, to give the product as a thick oil (0.32 g, 59%).

$^1\text{H-NMR}$ δ (CDCl_3) 1.23 (9H, d), 1.85 (8H, m), 2.58 (4H, brt), 3.24 (6H, m), 4.21 (2H, d), 4.37 (1H, m), 7.16 (4H, m), 7.23 (2H, m), 7.47 (1H, m), 8.42 (2H, brs).

5 Mass Spec. (ESI): 494.4 (MH) $^+$.

Example 2 N-Trimethylpyruvyl-L-pyrrolidinylmethyl-[N',N'-dibenzylamino]carbamate




10

$^1\text{H-NMR}$ δ (CDCl_3) 1.21 (s, 9H), 1.83 (m, 4H), 3.14 (m, 2H), 3.30 (m, 2H), 4.39 (brs, 4H), 4.48 (m, 3H), 7.28 (m, 10H).

Mass Spec. (ESI): 437.3 (MH) $^+$.

15

Example 3 N-Trimethylpyruvyl-L-pyrrolidinylmethyl-[N',N'-2-phenylethyl-benzylamino]carbamate




20

$^1\text{H-NMR}$ δ (CDCl_3) 1.23 (m, 9H), 1.91 (m, 4H), 2.80 (2 x t, 2H), 3.41 (m, 4H), 4.37 (m, 2H), 4.41 (brs, 1H), 7.21 (m, 10H).

Mass Spec. (ESI): 451.3 (MH) $^+$.


5 **Example 4** N-Trimethylpyruvyl-L-pyrrolidinylmethyl-[N'-3-pyridylmethylamino]carbamate



10 $^1\text{H-NMR}$ δ (CDCl_3) 1.24 (9H, s), 1.93 (4H, m), 3.35 (2H, m), 4.22 (2H, ABq), 4.39 (3H, d and m), 5.37 (1H, br), 7.32 (1H, m), 7.86 (1H, m), 8.55 (2H, m).

Mass Spec. (ESI): 379.3 (MH) $^+$.


15 **Example 5** N-Trimethylpyruvyl-L-pyrrolidinylmethyl-[N'-2-phenylethylamino]carbamate



20 $^1\text{H-NMR}$ δ (CDCl_3) 1.22 (s, 9H), 1.90 (m, 4H), 2.80 (t, 2H), 3.35 (m, 2H), 3.42 (q, 2H), 4.20 (d, 2H), 4.34 (m, 1H), 4.73 (m, 1H), 7.21 (m, 5H).

Mass Spec. (ESI): 361.4 (MH) $^+$.

Example 6 N-Trimethylpyruvyl-L-pyrrolidinylmethyl-[N'-3-phenylpropylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 1.21 (s, 9H), 1.82 (m, 6H), 2.60 (t, 2H), 3.16 (m, 2H), 3.32 (m, 2H), 4.19 (d, 2H), 4.32 (m, 1H), 4.96 (m, 1H), 7.20 (m, 5H).

Mass Spec. (ESI): 375.4 (MH) $^+$.

Example 7 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-phenylpropylamino]carbamate



15 A solution of N-t-butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-phenylpropylamino]carbamate (470.2 mg, 0.9763 mmoles) in methylene chloride (2 mL) was treated with 4M HCl in dioxane (2.5 mL). After 1 h the solvents were removed under vacuum. The residue was flushed with dry methylene chloride (3 x 25 mL), dried *in vacuo* for 1 h, and dissolved in dry methylene chloride (5 mL). To this


20 was added a solution of 3,4,5-trimethoxyphenyl-2-oxoacetyl chloride (506.5 mg, 2

equiv.) in methylene chloride (5 mL), followed by diisopropylethylamine (0.51 mL, 3 equiv.). The mixture was stirred at rt overnight and then concentrated *in vacuo*. The residue was dissolved in 1:1 ethyl acetate/ether and the solution washed with 1M sodium hydroxide (2x), water, brine, dried over sodium sulfate and evaporated. The residue was chromatographed on silica, eluting with ethyl acetate, to give the product as a pale-yellow oil (561.8 mg, 95%).

- ¹H-NMR δ (CDCl₃) 1.83 (6H, m), 1.94 (2H, m), 2.54 (4H, m), 3.03 (1H, m), 3.22 (3H, m), 3.41 (1H, m), 3.59 (1H, m), 3.84 (9H, m), 4.22 (1H, m), 4.37 (1H, m), 4.50 (1H, m), 7.17 (4H, m), 7.21 (4H, d and m), 7.44 (1H, brt), 8.40 (2H, brs).
- 10 Mass Spec. (ESI): 604.6 (MH)⁺.


The following compounds were prepared essentially as described for the previous example 7:

- 15 **Example 8 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N'-3-pyridylmethylamino]carbamate**



- 20 ¹H-NMR δ (CDCl₃) 1.96 (m, 4H), 3.52 (m, 2H), 3.81 (m, 9H), 4.00 (m, 2H), 4.30 (d, 2H), 4.41 (m, 1H), 5.80 (brt, 1H), 7.20 (m, 3H), 7.49 (ABq, 1H), 8.41 (m, 2H).
- Mass Spec. (ESI): 458.4 (MH)⁺.

Example 9 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N'-2-phenylethylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 1.96 (m, 4H), 2.58 (t, 2H, rotamer A), 2.79 (t, 2H, rotamer B), 3.00-3.80 (several m, 4H), 3.89 (m, 9H), 4.44 (m, 1H), 4.81 (brt, 1H), 7.19 (m, 7H).

Mass Spec. (ESI): 471.4 (MH) $^+$.

10

Example 10 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)butyl-4-phenylbutylamino]carbamate




15 $^1\text{H-NMR}$ δ (CDCl_3) 1.58 (8H, m), 1.95 (4H, m), 2.57 (4H, m), 2.90-3.70 (6H, 5 x m), 3.90 (9H, m), 4.32 (2H, ABm), 4.50 (1H, m), 7.17 (6H, m), 7.23 (2H, s), 7.42 (2H, d), 8.42 (2H, brs).

Mass Spec. (ESI): 632.4 (MH) $^+$.

36

Example 11 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-4-phenylbutylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 8.42 (d, 0.5 H); 8.39 (m, 0.5 H); 7.49-7.03 (m, 10 H); 4.49 (m, 3 H); 4.41-4.00 (m, 3 H); 3.90 (m, 9 H); 3.81-3.70 (m, 1 H); 3.63-3.02 (br m, 6 H); 2.57 (m, 2 H); 2.10-1.72 (m, 8 H).

Mass Spec. (ESI): 618.3 (MH) $^+$.

10

Example 12 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-2-phenylethylamino]carbamate




15

37

$^1\text{H-NMR}$ δ (CDCl_3) 8.42 (d, 0.5 H); 7.43 (m, 1 H); 7.35-7.02 (br m, 9 H); 4.59-4.20 (br m, 3 H); 3.92 (m, 9 H); 3.63 (m, 1 H); 3.41 (m, 2 H); 3.21 (m, 2 H); 3.01 (m, 0.5 H); 2.81 (m, 2 H); 2.5 (t, 0.5 H); 2.57 (m, 2 H); 2.15-1.45 (m, 6 H).
Mass Spec. (ESI): 590.3 (MH) $^+$.

5

Example 13 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-benzylamino]carbamate



10


$^1\text{H-NMR}$ δ (CDCl_3) 8.43 (m, 1 H); 7.43 (t, 1 H); 7.31-7.10 (m, 10 H); 4.55 (m, 0.5 H); 4.41 (m, 0.5 H); 4.24 (m, 1 H); 3.91 (m, 9 H); 3.51-2.98 (br m, 4 H); 2.59 (br s, 4 H); 2.05-1.71 (br m, 4 H); 1.55 (m, 2 H).
Mass Spec. (ESI): 576.3 (MH) $^+$.

15

20

38

Example 14 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)butyl-3-phenylpropylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 8.45 (m, 1.5 H); 7.51 (m, 0.5 H); 7.29-7.14 (m, 9 H); 4.59-4.21 (m, 3 H); 3.96 (s, 3 H); 3.90 (s, 3 H); 3.87 (s, 3 H); 3.43-2.95 (m, 6 H); 2.58 (m, 4 H); 2.05-1.67 (m, 6 H); 1.27 (br s, 4 H).

Mass Spec. (ESI): 618.4 (MH^+).

10

Example 15 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)butyl-2-phenylethylamino]carbamate



15


39

$^1\text{H-NMR}$ δ (CDCl_3) 8.41 (s, 1 H); 7.43 (m, 1 H); 7.21 (m, 5 H); 7.19 (m, 4 H); 4.52 (m, 1 H); 4.40 (m, 1 H); 4.23 (m, 1 H); 3.93 (s, 3 H); 3.91 (s, 6 H); 3.41 (m, 4 H); 3.20 (m, 2 H); 2.81 (q, 2 h); 2.62 (m, 2 H); 1.98 (m, 4 H); 1.57 (m, 4 H).

5 Mass Spec. (ESI): 604.3 (MH) $^+$.

**Example 16 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-
[N',N'-4-(3-pyridyl)butyl-benzylamino]
carbamate**

10




$^1\text{H-NMR}$ δ (CDCl_3) 8.41 (m, 1.5 H); 7.43 (m, 0.5 H); 7.30-7.05 (m, 9 H); 4.46 (m, 2 H); 4.37 (m, 1 H); 3.94 (s, 3 H); 3.90 (s, 6 H); 3.87 (s, 2 H); 3.47-3.11 (m, 15 4 H); 2.59 (m, 2 H); 2.10-1.43 (m, 8 H).

Mass Spec. (ESI): 590.3 (MH) $^+$.

20

40

Example 17 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-(3-methoxyphenyl)propylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 1.90 (m, 8H), 2.53 (m, 4H), 3.22 (m, 4H), 3.40 (m, 2H), 3.85 (m, 12H), 4.34 (m, 2H), 4.52 (m, 1H), 6.80 (m, 2H), 7.19 (t, 1H), 7.20 (ABq, 1H), 7.26 (m, 2H), 7.48 (t, 1H), 8.40 (m, 2H).

10 Mass Spec. (ESI): 634.43 (MH) $^+$.

Example 18 N-(3,4,5-Trimethoxyphenyl)oxalyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-(3,4,5-trimethoxyphenyl)propylamino]carbamate

15




41

$^1\text{H-NMR}$ δ (CDCl_3) 1.89 (m, 6H), 2.00 (m, 2H), 2.56 (2 x t, 4H), 3.24 (m, 4H), 3.41 (m, 2H), 3.87 (m, 18H), 4.30 (m, 2H), 4.53 (m, 1H), 6.39 (s, 2H), 7.21 (m, 3H), 7.46 (m, 1H), 8.41 (brs, 2H).

Mass Spec. (ESI): 694.45 (MH) $^+$.

5

Example 19 N-(3,4,5-Trimethoxyphenyl)oxalyl-(\pm)-piperidinylmethyl-[N',N'-3-(3-pyridyl)-propyl-3-phenylpropylamino]carbamate



10

$^1\text{H-NMR}$ δ (CDCl_3) 1.72 (m, 10H), 2.59 (m, 4H), 3.22 (m, 6H), 3.85 (s, 3H), 3.89 (s, 3H), 3.92 (s, 3H), 4.00-4.50 (m, 2H), 5.07 (m, 1H), 7.08 (s, 2H), 7.12 (m, 6H), 7.46 (d, 1H), 8.42 (brs, 2H).


Mass Spec. (ESI): 618.6 (MH) $^+$.

15

20

42

Example 20 N,N-(3-Phenylpropyl)-3-(3-pyridyl)propyl-2-[(S)-N'-(3,4,5-trimethoxyphenyl)oxalyl-2-piperidinyl]propionamide




5

$^1\text{H-NMR}$ δ (CDCl_3) 1.70 (m, 5 H), 1.86 (m, 6 H), 2.27 (m, 3 H), 2.61 (m, 4 H), 3.27 (m, 5 H), 3.85 (s, 3 H), 3.89 (s, 3 H), 3.93 (s, 3 H), 4.58 (m, 1 H), 4.82 (m, 1 H), 7.23 (m, 8 H), 7.52 (m, 1 H), 8.46 (m, 2 H).

Mass Spec. (ESI): 616 (MH) $^+$.

10

Example 21 N,N-(3-Phenylpropyl)-3-(3-pyridyl)propyl-2-[(S)-N'-(3,4,5-trimethoxyphenyl)oxalyl-2-pyrrolidinyl]propionamide




15

$^1\text{H-NMR}$ δ (CDCl_3) 1.85 (m, 11H), 2.38 (dt, 1 H, $J = 12.8, 7.7$), 2.60 (m, 4 H), 3.30 (m, 6 H), 3.95 (m, 9 H), 4.30 (m, 1 H), 7.22 (m, 8 H), 7.49 (m, 1 H), 8.44 (m, 2 H).

Mass Spec. (ESI): 602 (MH) $^+$.

5

Example 22 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)butyl-4-phenylbutylamino]carbamate



10

A solution of N-t-butyloxycarbonyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)butyl-4-phenylbutylamino]carbamate (50.7 mg, 0.0995 mmoles) in methylene chloride (1 mL) at rt was treated with 4M HCl in dioxane (0.25 mL, 10 equiv.). After 1 h the solvents were removed on the rotary evaporator at 30°C, and the residue flushed with dry methylene chloride (3 x 5 mL). The residue was dried *in vacuo* for 2 h, dissolved in dry methylene chloride, and then treated with 2,2-difluoro-3,4,5-trimethoxyphenylacetyl chloride (2 equiv.) and diisopropylethylamine (0.069 mL, 4 equiv.). After 12 h the mixture was concentrated *in vacuo* and the residue partitioned between ethyl acetate and 1M sodium hydroxide. The organic phase was washed with 1M sodium hydroxide, water and brine, dried over sodium sulfate, and evaporated.

20

The residue was chromatographed on silica, eluting with ethyl acetate, to give the product as a colorless gum (62.5 mg, 96%).

$^1\text{H-NMR}$ δ (CDCl_3) 1.55 (8H, m), 1.82 (4H, m), 2.60 (4H, m), 3.18 (4H, m), 3.43 (2H, m), 3.86 (9H, s), 4.22 (2H, m), 4.47 (1H, m), 6.79 (2H, s), 7.21 (6H, m),


5 7.48 (2H, d), 8.42 (2H, brs).

Mass Spec. (ESI): 654.4 (MH) $^+$.

The following compounds were prepared essentially as described for the previous example 22:

10

Example 23 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-phenylpropylamino]carbamate



15


$^1\text{H-NMR}$ δ (CDCl_3) 1.75 (m, 8H), 2.50 (m, 4H), 3.14 (m, 4H), 3.35 (m, 2H), 4.14 (m, 2H), 4.36 (m, 1H), 6.70 (s, 2H), 7.17 (m, 6H), 7.41 (m, 1H), 8.36 (m, 2H).

Mass Spec. (ESI): 626 (MH) $^+$.

20

45

Example 24 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-dibenzylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 7.38-7.09 (m, 10 H), 6.77 (s, 2 H), 4.50-4.28 (m, 7 H), 3.85(m, 9 H), 3.47 (m, 1 H), 3.25 (m, 1 H), 1.92-1.60 (m, 4 H).

Mass Spec. (ESI): 569.3 (MH) $^+$.

10 Example 25 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-2-(3-indolyl)ethyl-2-phenylethylamino]carbamate



15


46

$^1\text{H-NMR}$ δ (CDCl_3) 8.03 (bs, 1 H), 7.66-6.82 (m, 9 H), 6.79 (s, 2 H), 4.50-4.20 (m, 3 H), 3.87 (m, 9 H), 3.58-3.24 (m, 6 H), 2.98 (m, 1 H), 2.91 (m, 1 H), 2.82 (m, 1 H), 2.70 (m, 1 H), 1.97-1.70 (m, 4 H).

Mass Spec. (ESI): 636.4 (MH) $^+$.

5

Example 26 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-4-phenylbutylamino]carbamate



10


$^1\text{H-NMR}$ δ (CDCl_3) 8.42 (m, 1 H); 7.43 (t, 1 H); 7.33-7.11 (m, 7 H); 6.79 (s, 2 H); 4.41 (m, 1 H); 4.23 (m, 3 H); 3.85 (s, 9 H); 3.55-3.38 (m, 2 H); 3.34-3.07 (m, 3 H); 2.59 (m, 4 H); 1.97-1.63 (m, 6 H), 1.55 (m, 4 H).

15 Mass Spec. (ESI): 640.3 (MH) $^+$.

20

47

Example 27 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-2-phenylethylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 8.41 (m, 1 H); 7.47 (m, 1 H); 7.34-7.03 (m, 7 H); 6.73 (s, 2 H); 4.42 (m, 1 H); 4.23 (m, 2 H); 3.87 (s, 9 H); 3.57-3.04 (br m, 3 H); 2.79 (m, 2 H); 2.55 (m, 2 H) 2.00-1.69 (m, 9 H).

10 Mass Spec (ESI): 612.3 (MH^+).

Example 28 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-benzylamino]carbamate



15


48

$^1\text{H-NMR}$ δ (CDCl_3) 8.41 (d, 0.5 H); 8.39 (m, 0.5 H); 7.45-7.07 (br m, 8 H); 6.78 (s, 2 H); 4.56-4.21 (m, 4 H); 4.20-3.98 (br m, 1 H); 3.87 (s, 9 H); 3.58-3.12 (br m, 4 H); 2.59 (m, 2 H); 1.99-1.58 (br m, 6 H).

Mass Spec. (ESI): 598.3 (MH) $^+$.

5

Example 29 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)butyl-3-phenylpropylamino]carbamate



10


$^1\text{H-NMR}$ δ (CDCl_3) 8.45 (m, 1.5 H); 7.53 (m, 0.5 H); 7.28-7.16 (m, 7 H); 6.78 (s, 2 H); 4.23 (m, 1 H); 4.21 (m, 2 H); 3.87 (s, 9 H); 3.51-3.37 (m, 2 H); 3.25-3.05 (m, 4 H); 2.61 (m, 4 H); 1.96-1.72 (m, 6 H); 1.56 (m, 4 H).

15 Mass Spec. (ESI): 640.3 (MH) $^+$.

20

49

Example 30 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)butyl-2-phenylethylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 8.41 (s, 1 H); 7.43 (m, 1 H); 7.20 (m, 7 H); 6.79 (s, 2 H); 4.42 (m, 1 H); 4.22 (m, 2 H); 3.87 (s, 9 H); 3.41 (m, 4 H); 3.18 (m, 2 H); 2.79 (m, 2 H); 2.60 (m, 2 H); 1.82 (m, 4 H); 1.57 (m, 4 H).

10 Mass Spec. (ESI): 626.3 (MH^+); 606.3 (M-F^+).

Example 31 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-4-(3-pyridyl)butyl-benzylamino]carbamate



15


50

$^1\text{H-NMR}$ δ (CDCl_3) 8.41 (m, 1.5 H); 7.43 (m, 0.5 H); 7.40-7.13 (m, 7 H); 6.79 (s, 2 H); 4.78-4.21 (m, 3 H); 3.89 (s, 9 H), 3.86 (s, 2 H); 3.58-3.09 (m, 4 H), 2.59 (m, 2 H), 1.99-1.41 (m, 8 H).

Mass Spec. (ESI): 612.4 (MH^+).

5

Example 32 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-2-(3-methoxyphenyl)ethylamino]carbamate



10


$^1\text{H-NMR}$ δ (CDCl_3) 8.75 (m, 1 H), 8.63 (m, 1 H), 8.19 (m, 1 H), 7.80-7.28 (m, 5 H), 6.75 (s, 2 H), 4.50 (m, 1 H), 4.30-4.08 (m, 2 H), 3.86 (m, 12 H), 3.59-3.36 (m, 4 H), 3.32-3.10 (m, 2 H), 3.01-2.67 (m, 4 H), 2.02-1.69 (m, 6 H).

15 Mass Spec. (ESI): 642.4 (MH^+).

20

51

Example 33 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-2-(3,4-dimethoxyphenyl)ethylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 8.77 (m, 1 H), 8.68 (m, 1 H), 8.27-7.40 (m, 3 H), 6.82-6.63 (m, 5 H), 4.50 (m, 1 H), 4.28-4.13 (m, 2 H), 3.87 (m, 15 H), 3.54-3.31 (m, 4 H), 3.21 (m, 2 H), 2.84-2.68 (m, 4 H), 2.00-1.70 (m, 6 H).

10 Mass Spec. (ESI): 672.4 (MH) $^+$.

Example 34 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-2-(3-trifluoromethylphenyl)ethylamino]carbamate

15




52

$^1\text{H-NMR}$ δ (CDCl_3) 8.75 (m, 1 H), 8.68 (M, 1 H), 8.25-7.16 (m, 3 H), 6.82-6.65 (m, 5 H), 4.50 (m, 1 H), 4.30-4.13 (m, 2 H), 3.86 (m, 9 H), 3.55-3.33 (m, 4 H), 3.03 (m, 2 H), 2.88-2.66 (m, 4 H), 2.00-1.70 (m, 6 H).

Mass Spec. (ESI): 680.4 (MH) $^+$.

5

Example 35 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-(4-carboxyphenyl)propylamino]carbamate



10


$^1\text{H-NMR}$ δ (CDCl_3) 8.53-8.46 (m, 2 H), 8.02 (s, 1 H), 7.98 (s, 1 H), 7.72-7.19 (m, 4 H), 6.77 (s, 2 H), 4.47 (m, 1 H), 4.28-4.16 (m, 2 H), 3.86 (m, 9 H), 3.77-2.54 (m, 10 H), 1.98-1.72 (m, 8 H).

15 Mass Spec. (ESI): 670.4 (MH) $^+$.

20

53

Example 36 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-(4-benzyloxycarbonylphenyl)propylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 8.44 (m, 2 H), 8.00 (s, 1 H), 7.97 (s 1 H), 7.61-7.18 (m, 9 H), 6.78 (s, 2 H), 5.36 (s, 2 H), 4.45 (m, 1 H), 4.26-4.14 (m, 2 H), 3.87 (m, 9 H), 3.53-3.10 (m, 6 H), 2.70-2.51 (m, 4 H), 2.00-1.70 (m, 8 H).

10 Mass Spec. (ESI): 670.4 (MH) $^+$.

Example 37 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-(3-methoxyphenyl)propylamino]carbamate

15




54

$^1\text{H-NMR } \delta$ (CDCl_3) 1.81 (m, 8H), 2.55 (m, 4H), 3.21 (m, 4H), 3.43 (m, 2H), 3.78 (s, 3H), 3.85 (m, 9H), 4.21 (m, 2H), 4.42 (m, 1H), 6.73 (m, 3H), 6.77 (s, 2H), 7.20 (m, 2H), 7.47 (brt, 1H), 8.41 (brs, 2H).

Mass Spec. (ESI): 656.41 (MH^+).

5

Example 38 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-(4-methoxyphenyl)propylamino]carbamate



10


$^1\text{H-NMR } \delta$ (CDCl_3) 1.81 (m, 8H), 2.58 (m, 4H), 3.20 (m, 4H), 3.42 (m, 2H), 3.78 (s, 3H), 3.87 (m, 9H), 4.22 (ABq, 2H), 4.42 (m, 1H), 6.78 (s, 2H), 6.82 (d, 2H), 7.07 (m, 2H), 7.23 (ABq, 1H), 7.54 (m, 1H), 8.42 (brs, 2H).

15 Mass Spec. (ESI): 656.40 (MH^+).

20

55

Example 39 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-(3,4,5-trimethoxyphenyl)propylamino]carbamate



5

$^1\text{H-NMR}$ δ (CDCl_3) 1.82 (m, 8H), 2.58 (m, 4H), 3.21 (m, 4H), 3.43 (m, 2H), 3.84 (m, 18H), 4.20 (ABq, 2H), 4.42 (m, 1H), 6.37 (d, 2H), 6.77 (s, 2H), 7.21 (ABq, 1H), 7.48 (brt, 1H), 8.43 (m, 2H).


10 Mass Spec. (ESI): 716.47 (MH) $^+$.

15

20

56

Example 40 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-L-pyrrolidinylmethyl-[N',N'-3-(3-pyridyl)propyl-4-(3,4-dimethoxyphenyl)butylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 1.56 (m, 4H), 1.81 (m, 6H), 2.56 (m, 4H), 3.20 (m, 4H), 3.42 (m, 2H), 3.87 (m, 15H), 4.20 (m, 2H), 4.41 (m, 1H), 6.69 (brs, 2H), 6.78 (s, 2H), 6.79 (m, 1H), 7.22 (ABq, 1H), 7.50 (brt, 1H), 8.45 (m, 2H).

10 Mass Spec. (ESI): 700.49 (MH^+).

Example 41 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-(\pm)-piperidinylmethyl-[N',N'-3-(3-pyridyl)propyl-benzylamino]carbamate



15


57

$^1\text{H-NMR}$ δ (CDCl_3) 8.42 (d, 0.5 H); 8.40 (s, 0.5 H); 7.43 (m, 1 H); 7.21 (m, 7 h); 6.73 (d, 2 H); 5.07 (m, 1 H); 4.39 (m, 6 H); 3.85 (m, 9 H); 3.26 (t, 1 H); 3.20 (t, 1 H); 2.57 (m, 2 H); 1.84-1.35 (br m, 8 H).

Mass Spec. (ESI): 612.4 (MH^+); 592.4 (M-F^+).

5

Example 42 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-(\pm)-piperidinylmethyl-[N',N'-3-(3-pyridyl)propyl-2-phenylethylamino]carbamate



10


$^1\text{H-NMR}$ δ (CDCl_3) 8.41 (s, 1 H); 7.43 (m, 1 H); 7.21 (m, 7 H); 6.73 (d, 2 H); 5.07 (m, 1 H); 4.61-4.08 (br m, 3 H); 3.87 (s, 6 H); 3.83 (s, 3 H); 3.41 (m 2 H); 3.20 (m, 3 H); 2.82 (m, 2 H); 2.59 (m, 2 H); 1.83-1.05 (br m, 8 H).

15 Mass Spec.: 626.4 (MH^+); 606.4 (M-F^+).

20

58

Example 43 N-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-(±)-piperidinylmethyl-[N',N'-3-(3-pyridyl)propyl-3-phenylpropylamino]carbamate




5

$^1\text{H-NMR}$ δ (CDCl_3) 8.41 (s, 1 H); 7.51 (m, 1 H); 7.22 (m, 7 H); 6.73 (s, 2 H); 5.07 (m, 1 H); 4.60-4.09 (br m, 3 H); 3.87 (s, 6 H); 3.83 (s, 3 H); 3.74 (m, 1 H); 3.33-3.01 (br m, 2.5 H); 2.82 (m, 0.5 H); 2.61 (m, 2 H); 1.84-1.01 (br m, 15 H).

10 Mass Spec.: 654.5 (MH^+); 634.5 (M-F^+).

Example 44 N,N-(3-Phenylpropyl)-3-(3-pyridyl)propyl-2-[(S)-N'-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-2-piperidinyl]propionamide




15

$^1\text{H-NMR}$ δ (CDCl_3) 1.50 (m, 6 H), 1.83 (m, 5 H), 2.20 (m, 3 H), 2.60 (m, 4 H), 3.03 (m, 1 H), 3.21 (m, 2 H), 3.34 (m, 2 H), 3.68 (m, 1 H), 3.85 (s, 3 H), 3.86 (s, 3 H), 3.87 (s, 3 H), 4.76 (m, 1 H), 6.74 (m, 2 H), 7.24 (m, 6 H), 7.50 (m, 1 H), 8.44 (m, 2 H).

5 Mass Spec. (ESI): 638 (MH) $^+$.

Example 45 N,N-(3-Phenylpropyl)-3-(3-pyridyl)propyl-2-[(S)-N'-(3,4,5-Trimethoxyphenyl)-difluoroacetyl-2-pyrrolidinyl]propionamide



10

$^1\text{H-NMR}$ δ (CDCl_3) 1.75 (m, 10 H), 2.26 (m, 2 H), 2.59 (m, 4 H), 3.24 (m, 2 H), 3.34 (m, 2 H), 3.45 (m, 2 H), 3.85 (m, 9 H), 4.19 (m, 1 H), 6.67 (m, 2 H), 7.24 (m, 6 H), 7.50 (m, 1 H), 8.43 (m, 2 H).

15 Mass Spec. (ESI): 624 (MH) $^+$.

Example 46

FKBP12 Rotamase Inhibition Assay

The rotamase activity of FKBP-12 was measured by an adaptation of the assay described by Kofron et al.. (*Biochemistry*, 30, pp. 6127-6134 (1991)). The assay was carried out at 4°C with 1 mg chymotrypsin/mL of assay with succinyl-Ala-Leu-Pro-Phe-p-nitroanilide as the substrate. Chymotrypsin rapidly hydrolyzes the peptide

20

bond on the C-terminal side of the Phe of the *trans* form of the peptide and releases the chromogenic p-nitroaniline. The rate of the reaction is controlled by the rate of conversion of the *cis* form of the peptide to the *trans*-form, the reaction catalyzed by FKBP12. The apparent K_i values for inhibition of the rotamase activity were
5 determined by measuring decreases in the first order rate constant of the reaction catalyzed by FKBP12 as a function of the concentrations of the compounds described herein. K_i is the concentration of the compound that causes 50 percent inhibition of rotamase activity which is indicative of neurite outgrowth activity. The results are presented in Tables 1-4.

10

Example 47


Assay of Neurite Outgrowth in PC12 Cell Cultures

PC-12A rat pheochromocytoma cells are maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal calf serum and 5% calf serum at 37°C
15 and 5% CO₂. Cells to be assayed are plated at 10⁴ per well of a 24 well plate and allowed to attach for 4-18 h. The medium is then replaced with DMEM plus 0.1% BSA, submaximal concentrations of nerve growth factor (NGF) (as determined by neurite outgrowth assay), and varying concentrations of the FKBP12 binding compound (0.1nM-10μM) in a final concentration of 0.25% DMSO. Control cultures
20 are treated with NGF in the absence of the FKBP12 binding compound. After 72 h, cultures are fixed with 4% formalin in PBS, stained with Commassie Blue, and approximately 200 cells are counted in random fields of each well. Cells with neurites longer than one cell diameter are counted as a percentage of total number of cells.

25 The FKBP12 binding compounds of formula I utilized in this invention cause a significant increase in neurite outgrowth over control cultures.

Additionally, compounds of this invention may also show benefit as reversers of multidrug resistance (MDR) in cancer chemotherapy and as agents for the treatment of HIV infection. Nonimmunosuppressive compounds possessing the structural elements of the FKBP12 binding portion of FK506 have shown utility in reversing P-glycoprotein mediated MDR (U.A. Germann, et al., *Anti-Cancer Drugs*, 8, pp. 125-140 (1997)). In addition, there has been no direct correlation shown between rotamase inhibitory activity and MDR reversing activity (J.R. Hauske, et al., *Bioorg. Med. Chem. Lett.*, 4, pp. 2097-2102 (1994)). In the area of HIV infection, it is known that immunophilins, including the FK506 binding proteins (FKBPs), are involved in facilitating binding of the HIV envelope protein gp120 to host CD4 receptors (M.M. Endrich, et al., *Eur. J. Biochem.*, 252, pp. 441-446 (1998)), and that FK506 inhibits the growth of HIV-infected cells (A. Karpas, et al., *Proc. Natl. Acad. Sci USA*, 89, pp. 8351-8355 (1992)).

Table 1. FKBP12 rotamase inhibition data with selected examples.




Ex. #	R ₁	R ₂	R ₃	X	% inhib. at 10 uM or 1 uM*
1	3-phenylpropyl	3-(3-pyridyl)-propyl	t-butyl	O	93
4	H	3-pyridylmethyl	t-butyl	O	12
3	benzyl	2-phenylethyl	t-butyl	O	39
2	benzyl	benzyl	t-butyl	O	27
5	H	2-phenylethyl	t-butyl	O	0

8	H	3-pyridylmethyl	3,4,5,-trimethoxyphenyl	O	11
6	H	3-phenylpropyl	t-butyl	O	32
9	H	2-phenylethyl	3,4,5,-trimethoxyphenyl	O	11
25	2-phenylethyl	indolyethyl	3,4,5,-trimethoxyphenyl	F ₂	11*
24	Benzyl	benzyl	3,4,5,-trimethoxyphenyl	F ₂	34*

Table 2. FKBP12 rotamase inhibition data with selected examples.

5




Example #	y	X	n	m	% inhib. at 10 uM or 1uM*	K _i , nM
13	1	O	3	1	79	
28	1	F ₂	3	1	99	577
12	1	O	3	2	80	
27	1	F ₂	3	2	97	234
7	1	O	3	3	100	447
23	1	F ₂	3	3	98	113
11	1	O	3	4	96	537
26	1	F ₂	3	4	99	158
16	1	O	4	1	18*	
31	1	F ₂	4	1	63*	686

63

15	1	O	4	2	82	
30	1	F ₂	4	2	97	
14	1	O	4	3	54*	
29	1	F ₂	4	3	80*	
10	1	O	4	4	52*	
22	1	F ₂	4	4	83*	271
41	2	F ₂	3	1	11*	
42	2	F ₂	3	2	1*	
19	2	O	3	3	11	
43	2	F ₂	3	4	0*	

Table 3. FKBP12 rotamase inhibition data with selected examples.



5

Example #	R	X	% rotamase inhib. at 1 uM
32		F ₂	93
33		F ₂	96
34		F ₂	96

64










37		F ₂	93
17		O	75
38		F ₂	95
39		F ₂	92
18		O	68
35		F ₂	94
36		F ₂	77
40		F ₂	98

Table 4. FKBP12 rotamase inhibition data with selected examples.



5

10

15

Example #	n	R ₁	R ₂	X	% rotamase inhib. at 1 uM
20	2	3-phenylpropyl	3-(3-pyridyl)-propyl	O	0
44	2	3-phenylpropyl	3-(3-pyridyl)-propyl	F ₂	0
21	1	3-phenylpropyl	3-(3-pyridyl)-propyl	O	1
45	1	3-phenylpropyl	3-(3-pyridyl)-propyl	F ₂	42

If pharmaceutically acceptable salts of the compounds of formula I are used, those salts are preferably derived from inorganic or organic acids and bases. Included among such acid salts are the following: acetate, aspartate, bisulfate, butyrate, citrate, fumarate, hydrochloride, hydrobromide, hydroiodide, lactate, maleate, oxalate, persulfate, propionate, succinate, tartrate. Base salts include ammonium salts, alkali metal salts, such as sodium and potassium salts, alkaline earth metal salts, such as calcium and magnesium salts, salts with organic bases, such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth.

Pharmaceutically acceptable carriers that may be used in these pharmaceutical compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir.

It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of compound of formula I will also depend upon the particular FKBP12 binding compound in the composition.


10 The amount of compound of formula I utilized in these methods is between about 0.01 and 100 mg/kg body weight/day.

67

CLAIMS

What is claimed is:

- 5 1. A compound having the formula (I)



and pharmaceutically acceptable salts thereof, wherein:

10

Z is O, NH, N(C₁-C₃)-alkyl or CH₂;

X is O or F₂;

- 15 B and C are independently hydrogen, (C₁-C₄)-alkyl, or benzyl;

J is hydrogen, (C₁-C₄)-alkyl or benzyl;

- 20 K is (C₁-C₄)-straight or branched alkyl, benzyl or cyclohexylmethyl, or wherein J and K may be taken together to form a 4-7 membered heterocyclic ring which may contain a heteroatom selected from the group consisting of O, S, SO or SO₂, N, and NR, wherein R is selected from the group consisting of hydrogen, (C₁-C₄)-straight or branched alkyl, (C₂-C₄)-straight or branched alkenyl or alkynyl, and (C₁-C₄)-bridging alkyl wherein a bridge is formed between the nitrogen and a carbon atom of said

heteroatom-containing chain to form a ring, and wherein said ring is optionally fused to an Ar group;

wherein the stereochemistry at carbon position 1 is R or S;

- 5 D is (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl, (C₅-C₇)-cycloalkyl or (C₅-C₇)-cycloalkenyl substituted with (C₁-C₄)-straight or branched alkyl or (C₂-C₄)-straight or branched alkenyl, O-(C₁-C₄)-straight or branched alkyl, O-(C₂-C₄)-straight or branched alkenyl, 2-indolyl, 3-indolyl, [(C₁-C₄)-alkyl or (C₂-C₄)-alkenyl]-Ar or Ar;

10

- Ar is a carbocyclic aromatic group selected from the group consisting of phenyl, 1-naphthyl, 2-naphthyl, indenyl, azulenyl, fluorenyl, and anthracenyl; or a heterocyclic aromatic group selected from the group consisting of 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,3-triazolyl, 1,3,4-thiadiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-triazinyl, 1,3,5-trithianyl, indoliziny, indolyl, isoindolyl, 3H-indolyl, indolinyl, benzo[b]furanyl, benzo[b]thiophenyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, 4H-quinoliziny, quinolinyl, isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, and phenoxazinyl;
- 15
- 20

Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halogen, hydroxyl, hydroxymethyl, nitro,


- 25 trifluoromethyl, trifluoromethoxy, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl, O-[(C₁-C₄)-straight or branched alkyl], O-benzyl, O-phenyl, 1,2-methylenedioxy, amino, carboxyl, N-[(C₁-C₅)-straight or branched alkyl or (C₂-C₅)-straight or branched alkenyl] carboxamides, N,N-di-[(C₁-C₅)-straight or branched

alkyl or (C₂-C₅)-straight or branched alkenyl] carboxamides, N-morpholinecarboxamide, N-benzylcarboxamide, N-thiomorpholinocarboxamide, N-picolinoylcarboxamide, O-W, CH₂-(CH₂)_p-W, O-(CH₂)_p-W, (CH₂)_p-O-W, and CH=CH-W;

5

W is 4-methoxyphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrazyl, quinolyl, 3,5-dimethylisoxazolyl, isoxazolyl, 2-methylthiazoyl, thiazoyl, 2-thienyl, 3-thienyl, or pyrimidyl; p is 0-2;

- 10 Q and A are independently hydrogen, Ar, (C₁-C₁₀)-straight or branched alkyl, (C₂-C₁₀)-straight or branched alkenyl or alkynyl, provided the carbon atom having the triple bond in the alkynyl group is not directly bonded to the nitrogen atom of the core; (C₅-C₇)-cycloalkyl substituted (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, (C₅-C₇)-cycloalkenyl substituted
- 15 (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, or Ar-substituted (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl wherein, in each case, any one of the CH₂ groups of said alkyl, alkenyl or alkynyl chains may be optionally replaced by a heteroatom selected from the group consisting of O, S, SO, SO₂, N, and NR, wherein R is selected from the group
- 20 consisting of hydrogen, (C₁-C₄)-straight or branched alkyl, (C₂-C₄)-straight or branched alkenyl or alkynyl, and (C₁-C₄)-bridging alkyl wherein a bridge is formed between the nitrogen and a carbon atom of said heteroatom-containing chain to form a ring, and wherein said ring is optionally fused to an Ar group; or



25

G is hydrogen, (C₁-C₆)-straight or branched alkyl or (C₂-C₆)-straight or branched alkenyl or alkynyl; and

T is Ar or substituted 5-7 membered cycloalkyl with substituents at positions 3 and 4
5 which are independently selected from the group consisting of oxo, hydrogen, hydroxyl, O-(C₁-C₄)-alkyl, or O-(C₂-C₄)-alkenyl.

2. A compound of claim 1 wherein:

10 J and K are taken together to form a pyrrolidine ring;

the stereochemistry at carbon 1 is S;

B and C are each hydrogen;

X is F₂ or O;

Z is O or CH₂;

15 D is 3, 4, 5 trimethoxyphenyl or t-butyl;

A is 3-(3-pyridyl)propyl or 4-(3-pyridyl)butyl; and

Q is phenyl-substituted (C₁-C₆)alkyl, wherein phenyl is optionally substituted with one to three substituents independently selected from (C₁-C₆) alkyl, O-(C₁-C₆) alkyl, carboxyl and trifluoromethyl, wherein said alkyl is straight or branched.

20

3. A compound of claim 2 wherein Q is:

3-phenylpropyl;

3-(3,4,5-trimethoxyphenyl)propyl;

2-(3,4-dimethoxyphenyl)ethyl;

25 2-phenylethyl;

4-phenylbutyl;

4-(3,4-dimethoxyphenyl)butyl;

3-(4-carboxyphenyl)propyl;

2-(3-methoxyphenyl)ethyl;
2-(3-trifluoromethylphenyl)ethyl, or
3-(4-methoxyphenyl)propyl.

5 4. A pharmaceutical composition which comprises as an active ingredient an amount of a compound as claimed in any one of claims 1 to 3, or a pharmaceutically acceptable salt thereof, effective for stimulating neurite growth in nerve cells, and one or more pharmaceutically acceptable carriers, excipients or diluents thereof.

10 5. A method for stimulating neurite growth in nerve cells comprising the step of contacting said nerve cells with a composition comprising a neurotrophic amount of a compound with affinity for an FK-506 binding protein as claimed in any one of claims 1-3.

15 6. A method for stimulating neurite growth in nerve cells comprising the step of contacting said nerve cells with a composition comprising a neurotrophic amount of a compound with affinity for FKBP12 as claimed in any one of claims 1-3.

20

25

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/26798

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 207/08, 211/10; A61K 31/40, 31/445

US CL : 546/193; 548/518, 540; 514/318, 422, 423

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 546/193; 548/518, 540; 514/318, 422, 423

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

structure search: Caplus, Beilstein, Marpat

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,780,484 A (ZELLE et al) 14 July 1998(14.07.98).	1-6
A	WO 98/20891 A1 (ZELLE et al) 22 May 1998(22.05.98).	1-6

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

12 JANUARY 2000

Date of mailing of the international search report

23 FEB 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

LEIGH C. MAIER

Telephone No. (703) 308-1235